

清华大学深圳研究生院  
应用信息论  
2018 年春季学期

---

作业 1

YOUR NAME

2018 年 5 月 24 日

---

- 1.1. 设  $X$  和  $Y$  是各有均值  $m_x, m_y$ , 方差为  $\sigma_x^2, \sigma_y^2$ , 且相互独立的高斯随机变量, 已知  $U = X + Y, V = X - Y$ 。试求  $I(U; V)$ 。

解.  $U, V$  的联合分布是均值为  $[\mu_x + \mu_y, \mu_x - \mu_y]$ , 协方差矩阵为

$$\Lambda_{U,V} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \sigma_x^2 & 0 \\ 0 & \sigma_y^2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}^T = \begin{bmatrix} \sigma_x^2 + \sigma_y^2 & \sigma_x^2 - \sigma_y^2 \\ \sigma_x^2 - \sigma_y^2 & \sigma_x^2 + \sigma_y^2 \end{bmatrix}$$

由多元高斯分布微分熵的公式

$$h(U) = \frac{1}{2} \log((2\pi e)^2 |\Lambda_{U,V}|) = \frac{1}{2} \log(16\pi^2 e^2 \sigma_x^2 \sigma_y^2)$$

$U|V = v$  也是高斯分布, 方差为  $\frac{4\sigma_x^2 \sigma_y^2}{\sigma_x^2 + \sigma_y^2}$ , 与  $v$  无关, 因此

$$h(U|V) = \mathbb{E}_V[h(U|V = v)] = \frac{1}{2} \log(2\pi e \frac{4\sigma_x^2 \sigma_y^2}{\sigma_x^2 + \sigma_y^2}) \Rightarrow$$

$$\begin{aligned} I(U; V) &= h(U) - h(U|V) \\ &= \frac{1}{2} \log(16\pi^2 e^2 \sigma_x^2 \sigma_y^2) - \frac{1}{2} \log(2\pi e \frac{4\sigma_x^2 \sigma_y^2}{\sigma_x^2 + \sigma_y^2}) \\ &= \frac{1}{2} \log(2\pi e (\sigma_x^2 + \sigma_y^2)) \end{aligned}$$

- 1.2. 设有随机变量  $X, Y, Z$  均取值于  $\{0, 1\}$ , 已知

$$I(X; Y) = 0, I(X; Y|Z) = 1。求证 H(Z) = 1, H(X, Y, Z) = 2$$

证明.

$$I(X; Y|Z) = H(X|Z) - H(X|Y, Z) \leq H(X|Z) \leq H(X) \leq \log(2) = 1$$

所以等号全都成立  $\Rightarrow X \sim B(\frac{1}{2})$ 。同理可知  $Y \sim B(\frac{1}{2})$ 。另外

$$H(Y|Z) = H(Y) \Rightarrow I(Y; Z) = 0 \Rightarrow H(Z|Y) = H(Z)$$

$$\begin{aligned} H(X|Y, Z) &= 0 \\ \Leftrightarrow H(X, Y, Z) &= H(Y, Z) \\ \Leftrightarrow H(X, Y) + H(Z|X, Y) &= H(Y) + H(Z|Y) \\ \Leftrightarrow 2 + H(Z|X, Y) &= 1 + H(Z) \\ \Leftrightarrow H(Z) &= 1 + H(Z|X, Y) \end{aligned}$$

由上式推出  $H(Z) \geq 1$ , 又  
 $H(Z) \leq 1 \Rightarrow H(Z) = 1 \Rightarrow H(X, Y, Z) = 2$   $\square$

- 1.3. 设有信号  $X$  经过处理器  $A$  后获输出  $Y, Y$  再经处理器  $B$  后获输出  $Z$ 。  
已知处理器  $A$  和  $B$  分别独立处理  $X$  和  $Y$ 。试证:  $I(X; Z) \leq I(X; Y)$

**证明.**  $I(X; Z) = H(Z) - H(Z|X) = H(Z); I(Y; Z) = H(Y)$  因为  $Z$  是  $Y$  的函数  $\Rightarrow H(Z) \leq H(Y) \Rightarrow I(X; Z) \leq I(X; Y)$   $\square$

- 1.4. 已知随机变量  $X$  和  $Y$  的联合概率密度  $p(a_k, b_j)$  满足

$$p(a_1) = \frac{1}{2}, p(a_2) = p(a_3) = \frac{1}{4}, p(b_1) = \frac{2}{3}, p(b_2) = p(b_3) = \frac{1}{6}$$

试求能使  $H(X, Y)$  取得最大值的联合概率密度分布。

**解.**  $H(X, Y) = H(X) + H(Y) - I(X; Y) \leq H(X) + H(Y) = \frac{7}{6} + \log 3$   
等号成立当且仅当  $X, Y$  相互独立  $\Rightarrow p(x, y) = p(x)p(y)$

- 1.5. 设随机变量  $X, Y, Z$  满足  $p(x, y, z) = p(x)p(y|x)p(z|y)$ 。求证  
 $I(X; Y) \geq I(X; Y|Z)$

**证明.** 因为  $p(x, y, z) = p(x)p(y|x)p(z|y, x) \Rightarrow p(z|y, x) = p(z|x) \Rightarrow z$  与  $y$  条件独立  $\Rightarrow I(X; Y|Z) = H(X|Z) - H(X|Y, Z) = H(X|Z) - H(X|Y) \leq H(X) - H(X|Y) = I(X; Y)$   $\square$

- 1.6. 求证  $I(X; Y; Z) = H(X, Y, Z) - H(X) - H(Y) - H(Z) + I(X; Y) + I(Y; Z) + I(Z; X)$ ,  
其中  $I(X; Y; Z) \triangleq I(X; Y) - I(X; Y|Z)$

**证明.**

$$\begin{aligned} I(X; Y; Z) &= I(X; Y) - I(X; Y|Z) \\ &= H(X) + H(Y) - H(X, Y) - (H(X|Z) - H(X|Y, Z)) \\ &= H(X) + H(Y) - H(X, Y) - (H(X, Z) - H(Z)) + H(X, Y, Z) - H(Y, Z) \\ &= H(X, Y, Z) - H(X) - H(Y) - H(Z) + (H(X) + H(Y) - H(X, Y)) \\ &\quad + (H(Y) + H(Z) - H(Y, Z)) + (H(Z) + H(X) - H(X, Z)) \\ &= H(X, Y, Z) - H(X) - H(Y) - H(Z) + I(X; Y) + I(Y; Z) + I(Z; X) \end{aligned}$$

$\square$

1.7. 令  $p = (p_1, p_2, \dots, p_a)$  是一个概率分布, 满足  $p_1 \geq p_2 \geq \dots \geq p_a$ , 假设  $\epsilon > 0$  使得  $p_1 - \epsilon \geq p_2 + \epsilon$  成立, 证明:

$$H(p_1, p_2, \dots, p_a) \leq H(p_1 - \epsilon, p_2 + \epsilon, p_3, \dots, p_a)$$

**证明.** 设  $f(\epsilon) = (p_1 - \epsilon) \log(p_1 - \epsilon) + (p_2 + \epsilon) \log(p_2 + \epsilon)$  由已知

$$0 \leq \epsilon \frac{p_2 - p_1}{2} f'(\epsilon) = \log \frac{p_2 + \epsilon}{p_1 - \epsilon} \leq 0$$

$$\Rightarrow f(\epsilon) \leq f(0) \Rightarrow H(p_1, p_2, \dots, p_a) \leq H(p_1 - \epsilon, p_2 + \epsilon, p_3, \dots, p_a) \quad \square$$

1.8. 设  $p_i(x) \sim N(\mu_i, \sigma_i^2)$ , 试求相对熵  $D(p_2||p_1)$

**解.**

$$\begin{aligned} D(p_2||p_1) &= \int_{\mathbb{R}} p_2(x) \log \frac{p_2(x)}{p_1(x)} dx \\ &= \int_{\mathbb{R}} p_2(x) \left( \log \frac{\sigma_1^2}{\sigma_2^2} + \frac{1}{2}((x - \mu_1)^2 - (x - \mu_2)^2) \log e \right) dx \\ &= 2 \log \frac{\sigma_1}{\sigma_2} + \frac{1}{2}(\mu_1^2 - \mu_2^2) \log e + (\mu_2 - \mu_1)\mu_2 \log e \\ &= 2 \log \frac{\sigma_1}{\sigma_2} + \frac{1}{2}(\mu_1 - \mu_2)^2 \log e \end{aligned}$$

1.9. 若  $f(x)$  分别是区间  $(0, 0.01), (0, 0.5), (0, 1), (0, 2), (0, 5)$  上均匀分布的分布函数, 计算  $f(x)$  的微分熵。

**解.** 设  $U_t$  是  $(0, t)$  上的均匀分布, 则  $h(U_t) = \log t$

- $h(U_{0.01}) = \log 0.01$
- $h(U_{0.5}) = -1$
- $h(U_1) = 0$
- $h(U_2) = 1$
- $h(U_5) = \log 5$

1.10. 设

$$p_1(x, y) = \frac{1}{2\pi\sigma_x\sigma_y} \exp\left[-\frac{1}{2}\left(\frac{x^2}{\sigma_x^2} + \frac{y^2}{\sigma_y^2}\right)\right]$$

$$p_2(x, y) = \frac{1}{2\pi\sigma_x\sigma_y\sqrt{1-\rho^2}} \exp\left[-\frac{1}{2(1-\rho^2)}\left(\frac{x^2}{\sigma_x^2} - 2\rho\frac{xy}{\sigma_x\sigma_y} + \frac{y^2}{\sigma_y^2}\right)\right]$$

试求  $D(p_2||p_1)$  和  $I(X; Y)$ , 其中  $X, Y \sim p_2$

解.

$$\begin{aligned}
 D(p_2||p_1) &= \iint_{\mathbb{R}^2} p_2(x, y) \log \frac{p_2(x, y)}{p_1(x, y)} dx dy \\
 &\quad - \frac{1}{2} \log(1 - \rho^2) \\
 &\quad - \frac{1}{2} (\log e) \iint_{\mathbb{R}^2} p_2(x, y) \left[ \frac{\rho^2 x^2}{\sigma_x^2(1 - \rho^2)} + \frac{\rho^2 y^2}{\sigma_y^2(1 - \rho^2)} - \frac{2\rho xy}{(1 - \rho^2)\sigma_x\sigma_y} \right] dx dy \\
 &= -\frac{1}{2} \log(1 - \rho^2)
 \end{aligned}$$

$X|Y = y$  服从高斯分布, 方差为  $(1 - \rho^2)\sigma_x^2$

$$\begin{aligned}
 I(X; Y) &= h(X) - h(X|Y) \\
 &= \frac{1}{2} \log(2\pi e \sigma_x^2) - \frac{1}{2} \log(2\pi e \sigma_x^2 (1 - \rho^2)) \\
 &= \frac{1}{2} \log\left(\frac{2\pi e}{1 - \rho^2}\right)
 \end{aligned}$$